

## **ANTIWRINKLE COMPOSITION AND AGE REVERSAL COMPLEX**

[0001] This application claims the benefit of U.S. Provisional Application No. 60/410,555, filed August 8, 2002.

### **BACKGROUND OF THE INVENTION**

5 [0002] This invention relates to a cosmetic composition contain carnosine that is highly effective in controlling the ageing process by reversing, preventing and reducing wrinkling of skin. In particular the invention includes a complex containing carnosine that is especially effective in reversing, preventing and reducing the aging process that is visually observable through the formation of skin wrinkles. During the  
10 aging process glycated end products of bonding between protein material and sugar molecules naturally leads to formation of deep wrinkles in the skin that are particularly visible in normally exposed areas such as the face. Without being bound to the reasons for the effectiveness of the product of the present invention, it is theorized that carnosine prevents, reduces and can reverse the formation of such  
15 glycated end products and also breaks the protein-sugar bonds to enable actual reversal of skin wrinkles. If glycosylation were not prevented it would proceed as follows: a. sugar/aldehyde molecules approach protein molecules (for instance, malondialdehyde could approach, attach to and inactivate healthy protein molecules); b. sugar molecules attach to protein molecules leading to carbonylation crosslinking;  
20 c. the proteins that are carbonylated crosslinks (a detrimental factor); and d. advanced glycosylation end products (GRE's) form, which can cause extensive damage by reacting with free radicals and other toxins so that AGE's deposit in the cell structure (it is noted in this regard that malondialdehyde is the result of free radical damage to lipids, the fatty portion of cells, leading to a vicious cycle of aging (observed by skin  
25 wrinkling) since oxidation results in free radical formation that produces malondialdehyde, causing glycosylation that results in AGE's leading to more free radicals to repeat the process).

[0003] Prior research has determined that carnosine, with its known anti-glycating properties, may provide an anti-aging effect on fibroblast material by  
30 inhibiting cross-linking of glycoxidized proteins (Hipkiss et al, Mech. Ageing Dev., Sept. 15, 2000, 122(13), p. 1431-1445). Carnosine is reactive as an anti-glycating agent with aldehydes and ketones, that is compounds containing carbonyl groups (Hipkiss et al, Cell Mol. life Sci., May, 2000, 57 (5), p. 747-753 capable of blocking

carbonylated proteins, that form in step b., describe above, and thus prevent the formation of detrimental AGE's.

[0004] Carnosine and complexes containing carnosine have been described in skin care compositions and to promote wound healing, as in US Patents 5,091,171 to Yu et al; 5,723,482 to Degwert et al; 5,606,588 to Zaloga et al; and 4,981,846 to Matsukura et al. It is noteworthy, with particular regard to the patent to Yu et al, that while various cosmetic effects are disclosed for compositions containing an alpha-hydroxyacid or alpha-ketoacid and various amphoteric materials, including dipeptides. It is specifically disclosed that although the presence of alkali (ammonium hydroxide) does permit the composition to be therapeutic for dry skin, it loses much of its potency against wrinkles. Carnosine is disclosed among the dipeptides set forth. However,, the patent does not recognize that it could be effective against skin wrinkles under alkaline conditions.

[0005] It is an advantage of this invention that an alkaline skin cosmetic composition is provided that is topically applied and transdermally effective against wrinkling of skin.

[0006] It is a further advantage of this invention that the anti-wrinkle composition contains carnosine, derivatives thereof or mixtures of carnosine and derivatives thereof.

[0007] It is a further advantage of this invention that the carnosine and/or its derivatives are present in a complex that can be observed to reverse, prevent and reduce the aging process with regard to reversing, preventing and reducing formation of skin wrinkles.

[0008] Additional advantages resulting from the use of carnosine, including its derivatives include:

1. Ability to combine with sugar aldehydes such as malondialdehyde and inactivate them, thereby preventing carbonylation crosslinking;
2. As antiglycosylating agents they are superior in activity against free radicals to usual antioxidants, such as Vitamins E and C and the like that are not antiglycosylating agents;
3. They are antiglycosylating nutrients;
4. They inactivate free radicals within the cell after their formation;
5. They increase elimination of materials already damaged by free radicals;

6. They are effective against chronic inflammation;
7. They act as chelating agents to bind heavy metals and inactivate them, as well as assisting in regulating the concentration of these heavy metals (especially zinc and copper) to raise or lower the concentrations to desired levels;
8. They have antioxidant effect by stabilizing cell membranes and preventing free radical damage especially to the cell membranes;
9. They reduce or block excessive nitric oxide activity by blocking activation of guanylate cyclase, which is normally activated by nitric oxide.

10 [0009] It is a further advantage of this invention that the alkaline composition is compatible with other cosmetically desirable components, particularly non-allergenic components.

[0010] Additional advantages will be apparent from consideration of the following specification.

#### 15 **BRIEF DESCRIPTION OF THE INVENTION**

[0011] In accordance with certain of its aspects, this invention relates to an alkaline cosmetic composition having a pH of about 7.5 to about 9.5 that is effective against skin wrinkling, said composition comprising carnosine, derivatives thereof and mixtures of carnosine and derivatives thereof dispersed in an alkaline cosmetically acceptable vehicle in effective antiwrinkle amount up to about 50% by weight of carnosine

#### **DETAILED DESCRIPTION OF THE INVENTION**

[0012] Carnosine is a naturally occurring dipeptide composed of  $\beta$ -alanine and histidine and has a molecular weight of 226.24. It is commonly found in brain and innervated tissues and the ocular lens. Advantages obtained with the use of carnosine in the present invention are not characteristics of  $\beta$ -alanine and histidine, even when use in mixture with each other. It may be employed in its  $\alpha$ - and  $\beta$ -forms as well as mixtures thereof. The  $\alpha$ -form (also called L-carnosine) is preferred and the most readily available. It is most preferably used in its uncomplexed form although it may in a complex salt or salts, for instance with zinc, copper or manganese or in uncomplexed or such complex salts of derivative compounds, such as N-acetylcarnosine, which is a lipid soluble acid. When this lipid soluble acid is used, the pH range of about 7.5 to about 9.5 is maintained with use of alkaline water in the

composition or by buffers. Other derivatives of carnosine included in the present invention are anserine (that is methylation of carnosine), homocarnosine, N- $\alpha$ -acetylcarnosine, carcinine and numerous variants that are imidazole di-peptides. Such derivatives are described in the book, Carnosine And Other Elixirs of Youth-The  
5 Miraculous Anti-Ageing Supplement, Kyriazis, 2003, Watkins Publishing, London.  
(Carnosine, a derivative thereof or mixture of carnosine and derivative thereof is employed in the cosmetic composition in an effective antiwrinkle amount typically about 0.5% to about 50% that is about 5 mg/ml to about 500 mg/ml of composition), preferably about 1.5% to about 30% and most preferably about 10% to about 20% by  
10 weight

[0013] Additional ingredients that cooperate with and are compatible with carnosine to achieve nutrient balance and improve anti-wrinkling effectiveness include cascading antioxidants such as rosmarinic acid (that is, carnosic acid or carnosol embilica and additional polyphenol antioxidants, as well as anti-  
15 inflammatory materials such as beta-glucan, alkyl alcohol amines and the like. In this regard, it is noted that carnosine possesses antioxidant and anti-inflammatory properties.

[0014] It is also desirable to provide carnosine or its derivative either as such or as a complex salt and a complex composition that contains one or more of  
20 penetration enhancers of skin such as phosphatidyl, choline, serine and L-theanine, neutralizing agents for reactive radicals such as tocotrienols, especially in the gamma form,  $\gamma$ -tocopherol and mixed  $\alpha$ -,  $\beta$ - and  $\gamma$ -tocotrienols, anti-inflammatory agents such as glycerhizinate of licorice extract, cell stimulants such as glutathione and kinetin, growth promoting and bio-activating agents such as L-tyrosine and N-  
25 acetylcystine, and enhancers of the immune system such as clostrum and lactoferrin. Synergistic effects can be observed with the carnosine complex.

[0015] Typical concentrations for these compatible additives are about 0.1% to about 5.0% by weight of antioxidants, about 0.1% to about 10% by weight of anti-inflammatory, about 0.1% to about 10% by weight of skin penetrating agents, about  
30 0.1% to about 5% by weight of neutralizing agents of reactive radicals, about 0.1% to about 2% by weight of anti-inflammatory agents, about 0.1% to about 1% to about 50% by weight of each of cell stimulants, growth

promoting and bio-activating agents, and immune system enhancers. The carnosine compositions and the carnosine complexes can be employed in topically applied cosmetic compositions such as those applied to the skin in daytime, at nighttime, to tone the skin, to wash skin surfaces such as the face as well as periorbitally around the eyes.

[0016] Water, adjusted to a pH of about 7.5 to about 9.5, preferably about 8.0 to about 8.5, is a desirable vehicle for carnosine and other active components. Carnosine and other ingredients, if present, are dispersed in the alkaline aqueous vehicle under ambient conditions. The vehicle may desirably also contain a humectant such as glycerine or sorbitol, typically in amount of about 1% to about 10% by weight. The vehicle may also be an emulsion with a lipid component such as liquid paraffin, squalene, glycerol distearate and the like present in a typical amount of about 0.001% to about 50% by weight. The entire composition is readily prepared free of allergenic components. When carnosine or derivative thereof is present with water only the composition is a suspension.

[0017] Carnosine is particularly effective in preventing and reducing wrinkling of skin and may be effective in removing wrinkles as well, due to its excellent anti-glycation properties and ability to break crosslinks associated with wrinkled skin or to prevent them from forming. The effectiveness of carnosine as a wrinkle care material is quite unique to it in comparison to other amino acids including simple amino acids such as alanine and histidine and polypeptides including dipeptides such as glycylglycine and the like.

[0018] Carnosine and the various other ingredients described above, that may be present together with other additives that provide nutrients to the skin and for other purposes well known in the art, may be present for treating epidermis (surface) and dermis (deep) (layers such as those described for wrinkle-care products in US Patent 5,939,078 to Fujimura et al. the disclosure of which is incorporated herein by reference.

[0019] Such incorporated other additives typically may comprise about 0.001 to about 50% by weight of the cosmetic composition and may include:

humectants, ultra oily components, such as hydrocarbons, esters, high carbon fatty acids (e.g. stearic), high carbon alcohols (e.g. stearyl) and sphingosine derivatives from natural sources, typically in amount of about 0.005 to about 30% by weight; sterols, typically in amount of about 0.005 to about 30% by weight;

5 surfactants, especially nonionic surfactants, typically in amount of about 0.005 to about 30% by weight; water-soluble polyhydric alcohols, typically in amount of about 0.1 to about 25% by weight; powders, typically in amount of about 0.001 to about 50% by weight;

silicones, typically in amount of about 0.005 to about 30%; as well as minor amounts

10 of non-deteriorating salts, thickeners, pH regulators, violet absorbers, coloring agents, medicinal components, perfumes and the like. Other additives are included in the illustrative examples below.

[0020] The invention is illustrated by the following illustrative examples. All amounts and ranges are by in parts by weight unless otherwise indicated.

15 **[0021] Example 1**

The anti-wrinkling composition set forth below is prepared by dispersing the several ingredients with stirring in the alkaline aqueous vehicle.

	<u>Component</u>	<u>Parts</u>
	L-carnosine	25
20	Gamma-tocotreniol	5
	Licorice root	1
	Embilica	1
	L-theanine	10
	Glutathione	10
25	L-tyrosine	5
	L-acetylcysteine	2
	Clostrum	1
	Lactoferrin	1
	Deionized water (buffered to	Q.S. to 100
30		pH 8.0
	(buffered to with aminomethylpropanol, triethanolamine, monoisopropanolamine and dimethylmonoethanolamine)	

[0022] The above composition is topically applied to skin. It is effective to

35 reverse ageing by reducing wrinkles on wrinkled skin when applied to the skin in a regimen twice a day after for two weeks..

[0023] The pH of the above composition may be varied using alkaline water and varying the pH with the buffer in the range of 7.5 to 9.5.

**[0024] Example 2**

The composition of Example 1 is prepared except that N-acetylcarnosine replaces carnosine. Again, skin wrinkling is reduced.

5

**[0025] Example 3**

A suspension containing 35 parts of carnosine in water buffered at pH 7.5 to 8.0 is prepared and is effective to reduce skin wrinkles.

10

15

20

25

30

35

40

**[0025] Example 4**

The following compositions (A-G) are applied during daytime hours for anti-wrinkling effectiveness and include age-reversal complex ingredients.

Ingredient	<u>Parts</u>			
	<u>4A</u>	<u>4B</u>	<u>4C</u>	<u>4D</u>
Phosphatidyl Choline	0.20	0.50	2.00	0.20
Sodium Hyaluronate	0.50	0.10	0.10	0.10
Sugarcane Extract	0.10	0.10	0.10	0.10
EDTA-Na <sub>2</sub> *	0.10	-----	-----	0.10
Zinc Sulfate	-----	-----	-----	0.10
Na Carboxymethyl Betaglucan	0.10	-----	-----	-----
Xanthan Gum	-----	-----	-----	-----
Panthenol	0.05	-----	-----	-----
Et-hexyl Methoxycinnamate	7.00	7.00	7.00	7.00
Benzophenone-3	6.00	6.00	6.00	6.00
Isopropyl Palmitate	-----	-----	-----	5.00
Cetearyl Alcohol/ ceteaarth-20	-----	-----	-----	3.00
Cetearyl Alcohol 50/50	3.75	3.75	3.75	0.50
Glyceryl Stearate/ PEG* 100 Stearate	-----	-----	-----	1.00
Cetyl Esters	3.75	3.75	3.75	-----
Sorbitan Stearate	1.50	1.50	1.50	-----
Tetradecyl Ascorbate	0.10	0.25	0.50	0.10
Mg Boroascorbate-VDF	-----	-----	0.25	0.20
Vitamin A Palmitate	0.05	0.10	-----	-----
Tocotrienol R40 (Vitamin E)	0.10	0.20	0.50	0.10
Polysorbate 60	1.50	1.50	1.50	-----
Ubiquinone (Ubidecarenone)	0.01	-----	-----	-----
Super Oxide Dismutase	0.05	-----	-----	-----
PEG*-40 Castor Oil	0.25	1.50	1.50	-----
Squalane	1.00	-----	-----	-----
Benzyl Alcohol	1.50	1.50	1.50	-----
Phenoxyethanol & Me, Et, But, Pro & Isobut Parabens	-----	-----	-----	0.25
Rosemary Extract	-----	0.20	0.50	0.20
L-Carnosine	15.00	15.00	15.00	15.00
L-Theanine	2.50	2.50	-----	-----
Green Tea Extract-VDF	0.05	-----	-----	-----
Grape Seed Extract-VDF	0.05	-----	-----	-----
Vitamin K (powder)	0.05	0.10	0.10	0.10
Colostrum (powder)	0.05	0.05	0.05	0.05
Emblica (powder)	0.10	0.05	0.05	0.05
Licorice (powder)	-----	-----	-----	-----
Mn Gluconate*2 H <sub>2</sub> O	0.05	0.05	0.05	0.05
N-Acetyl L-Cysteine	-----	0.01	0.01	0.01
Dimethylethanolamine	0.10	0.15	0.15	0.15
Bergamot Oil	-----	0.001	0.001	0.001
Water (@ pH 8.0) q. s.100		100	100	100

\*EDTA Na<sub>2</sub> is disodium ethylene tetraacetic acid/PEG is polyethylene glycol

<u>Ingredient</u>	<u>Parts</u>		
	<u>4E</u>	<u>4F</u>	<u>4G</u>
Soy Lecithin	-----	-----	0.30
Phosphatidyl Choline	0.10	0.10	0.20
Sodium Hyaluronate	4.00	4.00	4.00
EDTA-Na <sub>2</sub>	0.10	0.10	0.10
Zinc Sulfate	-----	-----	0.10
Na Carboxymethyl Betaglucon	-----	-----	0.20
Xanthan Gum	-----	-----	0.10
Et-hexyl Methoxycinnamate	7.00	7.00	7.50
Benzophenone-3	6.00	6.00	6.00
Isopropyl Palmitate	5.00	5.00	-----
Cetearyl Alcohol/ceteaarth-20	3.00	3.00	-----
Cetearyl Alcohol 50/50	0.50	0.50	3.75
Glyceryl Stearate	-----	-----	-----
/PEG 100 Stearate	1.00	1.00	-----
Cetyl Esters	-----	-----	3.75
Sorbitan Stearate	-----	-----	1.50
Tetradecyl Ascorbate	0.25	0.25	0.10
Mg Boroascorbate-VDF	0.20	0.20	-----
Vitamin A Palmitate	-----	-----	0.10
Tocotrienol R40	0.50	0.50	0.10
Polysorbate 60	-----	-----	1.50
Squalane	-----	-----	0.20
Benzyl Alcohol	-----	-----	1.50
Phenoxyethanol & Me, Et, But, Pro & Isobut Parabens	0.30	0.30	-----
Rosemary Extract	0.30	0.30	-----
L-Carnosine	15.00	5.00	15.00
L-Theanine	-----	-----	2.50
Vitamin K (powder)	0.10	0.10	-----
Colostrum (powder)	0.05	0.05	0.30
Emblica (powder)	0.05	0.05	0.10
Licorice (powder)	-----	-----	0.10
Manganese Gluconate•2 H <sub>2</sub> O	0.05	0.05	-----
N-Acetyl L-Cysteine	0.01	-----	-----
Dimethylethanolamine	0.15	0.15	0.40
Bergamot Oil	0.001	0.001	-----
Water (@ pH 8.0)	q.s. 100	100	100

\*EDTA Na<sub>2</sub> is disodium ethylene diamine tetraacetic acid/PEG is polyethylene glycol

**[0026]** An aqueous pre-mix is prepared using Water, Phosphatidyl Choline and Sodium Hyaluronate as its base. Normally this aqueous pre-mix is heated to approximately 60° C to facilitate the dissolution of Phosphatidyl Choline. Other components of this premix—Soy Lecithin, Sugarcane Extract, EDTA-Na<sub>2</sub>, Zinc Sulfate, Na Carboxymethyl Betaglukan, Xanthan Gum, Emblica and Panthenol — are added in different combinations to vary the characteristics of the aqueous phase and final product.

**[0027]** A non-aqueous pre-mix is also prepared with Ethylhexyl Methoxycinnamate, Benzophenone-3, Cetearyl Alcohol 50/50, Tetrahexyldecyl Ascorbate and Tocotrienol R40 as its base. Normally this non-aqueous pre-mix is heated until all components are melted and dispersed throughout the pre-mix (~60° C). Isopropyl Palmitate, Cetyl Alcohol/Ceteareth-20, Glyceryl Stearate/PEG 100 Stearate, Cetyl Esters, Sorbitan Stearate, Vitamin A Palmitate, Squalane, Mg Ascorbate, Polysorbate 60, Ubiquinone (Ubidecarenone). Super Oxide Dismutase and PEG-40 Castor Oil are added in different combinations to vary the characteristics of the non-aqueous phase and final product.

**[0028]** After these two pre-mixes are prepared and heated, they are combined and the resulting mixture is stirred in order to achieve complete dispersion. It is then allowed to cool. Either Benzyl Alcohol or Phenoxyethanol & Me, Et, But. Pro & Isobut Parabens, used as a preservative system, is added to the cooling dispersion before the addition of L-Carnosine. To the resulting cooling dispersion, Vitamin K (powder), Colostrum (powder), Emblica (powder) and Manganese Gluconate•2 H<sub>2</sub>O are added in various combinations, as is Dimethylethanolamine, that is DMEA (to adjust the pH), along with the other remaining chosen ingredients (L-Theanine, Licorice Powder, Green Tea Extract – VDF, Grape Seed Extract – VDF, N-Acetyl L-Cysteine, Lavender Oil Fragrance and Bergamot Oil).

**[0029]** Upon reaching room temperature, the resulting cream is packaged

**[0030]** It is desirably applied one or two times a day over a two week period.

**[0031] Exmpl 5**

The following compositions(A-C) are applied during nighttime hours for anti-wrinkling effectiveness and include age-reversal complex ingredients (A and B are the most active; C is a simpler formula).

<u>Ingredient</u>	<u>Parts</u>		
	<u>5A</u>	<u>5B</u>	<u>5C</u>
Sodium Hyaluronate	0.10	----	----
Hydrolyzed Glycosaminoglycans	0.10	----	----
Sugarcane Extract	0.05	----	----
EDTA-Na <sub>2</sub>	0.10	0.10	0.10
Phosphatidyl Choline	----	4.00	----
Na carboxymethyl Betaglukan	0.05	----	0.05
Isopropyl Palmitate	6.25	6.25	6.25
Rice Bran Oil	----	1.00	----
Cetearyl Alcohol/Ceteareth-20 mix	3.75	3.75	3.75
Cetearyl Alcohol 50/50	0.60	0.60	0.60
Glyceryl Stearate/PEG 100 Stearate	1.25	1.25	1.25
Tetrahexyldecyl Ascorbate	0.10	0.10	0.50
Vitamin A Palmitate	0.05	0.10	0.01
Butylated Hydroxytoluene	0.30	----	0.30
PEG-40 Hydrogenated Castor Oil	0.25	0.25	1.00
Tocotrienol R40 (Vitamin E)	0.05	0.10	----
Ubiquinone (Ubidecarenone)	0.05	----	----
Super Oxide Dismutase	0.05	0.05	----
Mg Boroascorbate (VDF)	----	0.10	----
Squalane	1.00	----	----
Phenoxyethanol & Me, Et, but, Pro. & Isobut Parabens	0.20	0.20	0.20
L-Carnosine	15.00	15.00	15.00
L-Theanine	1.00	----	1.00
Green Tea Extract – VDF	0.05	0.05	----
Aloe Vera (200:1) Extract (Powder)	0.05	----	----
Colostrum (powder)	0.05	0.05	----
Emblica (powder)	0.10	0.10	----
Manganese Gluconate•2 H <sub>2</sub> O	0.05	0.05	----
Dimethylethanolamine	0.10	0.20	0.20
Bergamot Oil	----	----	<0.01
Water @pH 8.0	q.s. 100	100	100

**[0032]** An aqueous pre-mix is prepared using Water and EDTA-Na<sub>2</sub> (Na Carboxymethyl Betaglukan, Phosphatidyl Choline, Sugarcane Extract, Sodium Hyaluronate and Hydrolyzed Glycosaminoglycans are added in different combinations during the heating phase.) This aqueous pre-mix is heated to approximately 60° C.

**[0033]** A non-aqueous pre-mix is also prepared using Isopropyl Palmitate, Cetearyl Alcohol/Ceteareth-20, Cetearyl Alcohol 50/50, Glyceryl Stearate/PEG 100 Stearate, Tetrahexyldecyl Ascorbate, Vitamin A Palmitate and PEG-40 Hydrogenated Castor Oil as its base. Normally this non-aqueous pre-mix was heated until all components are melted and dispersed throughout the pre-mix (at approximately 60° C). For the 'active' non-aqueous premix, Squalane, Tocotrienol R40, Ubiquinone (Ubidecarenone), Superoxide Dismutase, Butylated Hydroxytoluene and PEG-40 Castor Oil (hydrogenated) are also added in various combinations to the mixture.

**[0034]** After the aqueous and non-aqueous pre-mixes are prepared and heated, they are combined and the resulting mixture is stirred in order to achieve complete dispersion. It is then allowed to cool. Phenoxyethanol & Me, Et, But, Pro & Isobut Parabens, used as a preservative system, is added to the cooling dispersion before the addition of L-Carnosine and L-Theanine. Dimethylethanolamine (DMEA) is added to adjust the pH in both dispersions, although it is added after the other remaining ingredients in the formulation (Green Tea Extract – VDF, Aloe Vera (200:1) Extract (Powder), Colostrum (powder), Emblica (powder), Mg Boroascorbate (VDF) and Manganese Gluconate•2 H<sub>2</sub>O in varying combinations). As a final step in the cooling 'simple' dispersion Bergamot Oil is added.

**[0035]** Upon reaching room temperature, the resulting cream was packaged. It is desirably applied at night over a two week period.

**[0036] Example 6**

The following compositions (A & B) are applied to tone the skin while providing anti-wrinkling effectiveness and included age-reversal complex ingredients.

<u>Ingredient</u>	<u>Parts</u>	
	A	B
L-Carnosine	1.00	5.00
L-Theanine	0.10	-----
Allantoin	0.05	0.10
Na PCA (Pyrocarboxylic Acid Na)	0.10	0.25
Sorbitol (70%)	0.25	0.10
Aloe Vera (200:1) Powder	0.05	0.01
Actiphyte of Witch Hazel	1.00	0.50
Actiphyte of Echinacea	1.00	0.50
Actiphyte of Chamomile	1.00	0.50
EDTA-Na <sub>2</sub>	0.10	0.05
PEG-60 Castor Oil (hydr)	0.05	0.05
Bergamot Oil	0.05	0.01
Phenoxyethanol & Me, Et, Bu, Pro, & Isobut Parabens	-----	0.10
Pentylene Glycol	-----	0.10
Water (@ pH 8.0)	<i>q.s.</i> 100	100

**[0037]** To the water, which is at room temperature, is added each of the other ingredients, one at a time, allowing for each to dissolve before the addition of the next. The additions are done while the aqueous solution is being stirred.

**[0038]** After sufficient stirring to insure a uniform solution, the resulting toner is packaged. It is desirably applied one to two times a day over a two week period.

**{0039} Example 7**

The following composition is applied to the face as a soft facial wash with anti-wrinkling effectiveness and includes age-reversal complex ingredients.

<u>Ingredient</u>	<u>Parts</u>
Mackadet EQ-76	
(Deionized water & PEG 80 Sorbitan	
Laurate, Cocamidopropyl	
Na Tridecath Sulfate, Na betaine	
Lauroamphoacetate, PEG 150	
Distearate, Na lauroeth S Carboxylate	
PEG 150 Distearate, Citric acid,	
Quaternium 15 & Na <sub>4</sub> EDTA)	52.00
L-Carnosine	5.00
Na PCA (Pyrocarboxylic Acid Na)	0.10
Jojoba Oil (Boston Jojoba)	0.01
Sugarcane Extract	0.25
Dimethanolethanolamine	0.15
Pomelo 7713 (Fragrance)	0.15
DMDM Hydantoin	0.15
Water	q.s. 100

**[0040]** The water is heated to ~ 45° C and the following ingredients are added one at a time to the water, with stirring, allowing enough time for dissolution of the previous ingredient: Mackadet EQ-76, L-Carnosine, Na PCA, Jojoba Oil (Boston Jojoba) and Sugarcane Extract. The aqueous solution is cooled to room temperature at which the remaining ingredients (Dimethylethanolamine, that is DMAE), Pomelo 7713 and DMDM Hydantoin) are added with stirring, allowing enough time for dissolution of the previous ingredient.

**[0041]** After sufficient stirring to insure a uniform solution, the resulting skin wash is packaged. It is desirably applied to wash the face once a day over a period of two weeks.

**[0042] Example 8**

The following compositions (A & B) are applied periorbitally around the eyes for their anti-wrinkling and age-reversal effects, preferably once daily over a two week period..

<u>Ingredient</u>	<u>Parts</u>	
	<u>8A</u>	<u>8B</u>
Phosphatidyl Choline	0.20	1.00
Sodium Hyaluronate	4.00	0.50
EDTA-Na <sub>2</sub>	0.10	0.10
Panthenol	0.10	0.10
Mg Boroascorbate-VDF	-----	0.25
Ethylhexyl Methoxycinnamate	5.00	5.00
Cetearyl Alcohol 50/50	3.75	3.75
Cetyl Esters	3.75	3.75
Sorbitan Stearate	1.50	1.50
Polysorbate 60	1.50	1.50
Squalane	0.10	-----
Ceramide- III	0.10	0.10
Tocotrienol R40 (Vitamin E)	0.10	0.20
Ubiquinone (Ubidecarenone)	0.10	0.10
Super Oxide Dismutase	0.10	0.10
Vitamin A Palmitate	0.05	0.05
Tetrahexyldecyl Ascorbate	0.10	0.10
Benzyl Alcohol	1.50	1.50
L-Carnosine	10.00	10.00
L-Theanine	0.50	0.50
Vitamin K (powder)	0.05	0.05
Green Tea Extract – VDF	0.05	0.05
Colostrum (powder)	0.10	0.10
Cucumber Extract	0.10	0.10
<u>Dimethylethanolamine</u>	<u>0.15</u>	<u>0.20</u>
Water (@ pH 8.0)	<i>q.s.</i> 100	100

**[0043]** An aqueous pre-mix is prepared using Water, Phosphatidyl Choline, Sodium Hyaluronate, EDTA-Na<sub>2</sub> and Panthenol as its base. Normally this aqueous pre-mix is heated to at approximately 60° C to facilitate the dissolution of Phosphatidyl Choline. Mg Boroascorbate is added to one of the formulations but not to the other.

**[0044]** A non-aqueous pre-mix was also prepared with Ethylhexyl Methoxycinnamate, Cetearyl Alcohol 50/50, Cetyl Esters, Sorbitan Stearate, Polysorbate 60, Ceramide III, Tocotrienol R40, Ubiquinone (Ubidecarenone), Super

Oxide Dismutase, Vitamin A Palmitate and Tetrahexyldecyl Ascorbate as its base. This non-aqueous pre-mix is heated until all components are melted and dispersed throughout the pre-mix (at approximately 60° C). Squalane is added to the pre-mix non-aqueous formulation for the product not containing Mg Boroascorbate in its aqueous pre-mix.

**[0045]** After these two pre-mixes are prepared, heated and combined and the resulting mixture is stirred in order to achieve complete dispersion. It is then allowed to cool. Benzyl Alcohol, used as a preservative system, is added to the cooling dispersion before the addition of L-Carnosine and L-Theanine. To the resulting cooling dispersion, Vitamin K (powder), Colostrum (powder), Green Tea Extract-VDF and Cucumber are added. The last step is the addition of Dimethylethanolamine (DMEA) to adjust the pH.

**[0046]** Upon reaching room temperature, the resulting cream is packaged.

**[0047]** It will be apparent that modifications and variations within the scope of the present invention may be prepared and provided.